

pay
as little as
\$5*

** Insured patients pay as little as \$5 co-pay per prescription. Maximum value of \$100 for 30-day prescriptions. See eligibility and restrictions.*



AMBRISENTAN TABLETS PATIENT SAVINGS CARD

Claims Processor: SimpleSaveRx

BIN # 017290

Rx PCN # 55101202

Group # X11020

Person Code: 01

Cardholder ID # 110203001001

PHARMACIST: please restore patient profile to primary PBM after claim submission.

For important safety information, see reverse side.

Complete prescribing information, including boxed warning, can be found on the Zydus website at www.zydususa.com.

PS-Ambrisentan REMS Program information can be found at <https://www.PSAmbrisentanREMS.com>.

DEAR PHARMACIST: SimpleSaveRx has been authorized to reimburse you per your contracted rate plus a maximum benefit determined by Zydus Pharmaceuticals for processing this card when accompanied by a valid prescription for Ambrisentan. Insured patients will receive a discount off their normal pharmacy charges if their co-pay exceeds \$5.

This claim may be submitted electronically through SimpleSaveRx using the processing numbers on the front of this co-pay card or by mail. Submit all electronic claims in NCPDP Standard D.0. Secondary processing should follow NCPDP standards for Co-Pay Only billing (other coverage code 3 or 8) or by using Coordination of Benefits processing.

Mail claims should go to SimpleSaveRx, 3350 N Arizona Ave, Ste. 2, Chandler, AZ 85225 along with the copy of the pharmacy prescription receipt (cash register receipts not accepted), and the return address. Retain a copy of this co-pay card and file with the prescription for auditing purposes. For expedited processing, Fax savings card and Rx receipt to: 480-444-1449.

TERMS & CONDITIONS: With this ambrisentan patient savings card, eligible patients can receive their Zydus Pharmaceuticals generic ambrisentan tablets 5 or 10 mg for as little as \$5 to a maximum value of \$100 per 30-day supply. The ambrisentan patient savings card is not health insurance.

ELIGIBILITY REQUIREMENTS: The ambrisentan patient savings card can only be used by eligible residents of the U.S., Puerto Rico, or U.S. Territories via Certified Pharmacies who dispense ambrisentan. Product must originate in the U.S. or Puerto Rico, or U.S. Territories. You must be 18 years or older to use the ambrisentan patient savings card.

The ambrisentan patient savings card is not insurance and is not intended to substitute for insurance.

Patients are eligible for this coupon savings program if they present a valid prescription for ambrisentan tablets and if they pay through commercial insurance. This offer may not be redeemed for cash and is not insurance. Where third-par-

ty reimbursement covers a portion of your prescription, this coupon is valid only for the amount of your actual out-of-pocket expenses, up to the maximum benefit allowed per this program.

The ambrisentan patient savings card is valid only for patients with commercial insurance and is not valid for prescriptions that are eligible to be reimbursed in whole or in part by Medicare, Medicare Part D, Medicaid (including Medicaid managed care), Medigap, Tricare, VA, DOD, FEHB, Puerto Rico Government Health Insurance Plan ("Healthcare Reform"), or any other federal or state-funded healthcare benefit program (collectively "Government Programs"); or by commercial plans or other health or pharmacy benefit programs that reimburse for the entire cost of the prescription drugs.

Medicare Part D Enrollees who are in the prescription drug coverage gap (the "donut hole") are not eligible for the ambrisentan patient savings card. Patients who begin receiving prescription benefits from Government Programs at any time will no longer be eligible to use the ambrisentan patient savings card. Void where prohibited by law, taxed, or restricted.

Patient, pharmacist, and prescriber agree not to seek reimbursement for all or any part of the benefit received by the patient through the offer. Both patient and pharmacist are each individually responsible for reporting receipt of the ambrisentan patient savings card benefit to any insurer, health plan, or other third party who pays for, or reimburses, any part of the prescription filled using the ambrisentan patient savings card, as required. It is illegal to sell, purchase, trade, or counterfeit, or offer to sell, purchase, trade, or counterfeit the ambrisentan patient savings card.

This offer is not transferrable. No substitutions are permitted. The offer cannot be combined with any other coupon, free trial, discount, prescription savings card, or other offer.

This offer is valid for Zydus Pharmaceuticals generic ambrisentan tablets 5 and 10mg only.

Call the SimpleSaveRx Help Desk at 1-844-SAVE4RX (844-728-3479) for processing questions.

Important Safety Information and Indication

Ambrisentan tablets are indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I):

- To improve exercise ability and delay clinical worsening. Studies establishing effectiveness included predominantly patients with WHO Functional Class II–III symptoms and etiologies of idiopathic or heritable PAH (60%) or PAH associated with connective tissue diseases (34%).

What is the most important information I should know about ambrisentan tablets?

Serious birth defects. Ambrisentan tablets can cause serious birth defects if taken during pregnancy.

- Females must not be pregnant when they start taking ambrisentan tablets or become pregnant during treatment with ambrisentan tablets.
- Females who are able to get pregnant must have a negative pregnancy test before beginning treatment with ambrisentan tablets, each month during treatment with ambrisentan tablets, and one month after stopping ambrisentan tablets. Talk to your doctor about your menstrual cycle. Your doctor will decide when to do a pregnancy test, and will order a pregnancy test for you depending on your menstrual cycle.

§ Females who are able to get pregnant are females who:

- have entered puberty, even if they have not started their menstrual period, and
- have a uterus, and
- have not gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed.

§ Females who are not able to get pregnant are females who:

- have not yet entered puberty, or
- do not have a uterus, or
- have gone through menopause. Menopause means that you have not had a menstrual period for at least 12 months for natural reasons, or that you have had your ovaries removed, or
- who are infertile for any other medical reason and this infertility is permanent and cannot be reversed

Females who are able to get pregnant must use two acceptable forms of birth control during treatment with ambrisentan tablets, and for one month after stopping ambrisentan tablets because the medicine may still be in the body.

- If you have had a tubal sterilization or have an IUD (intrauterine device) or progesterone implant, these methods can be used alone and no other form of birth control is needed.
- Talk with your doctor or gynecologist (a doctor who specializes in female reproduction) to find out about options for acceptable forms of birth control that you may use to prevent pregnancy during treatment with ambrisentan tablets.
- If you decide that you want to change the form of birth control that you use, talk with your doctor or gynecologist to be sure that you choose another acceptable form of birth control.

See the chart below for Acceptable Birth Control Options during treatment with ambrisentan tablets.

Acceptable Birth Control Options

Option 1	OR	Option 2	OR	Option 3	OR	Option 4
One method from this list:		One method from this list:		One method from this list:		One method from this list:
Standard intrauterine device (Copper T 380A IUD) Intrauterine system (LMg 20 IUS - Progesterone IUD) Tubal sterilization Progesterone implant		Estrogen and Progesterone oral contraceptives ("the pill") Estrogen and Progesterone transdermal patch Vaginal ring Progesterone injection		One method from this list: Diaphragm with spermicides Cervical cap with spermicides		Partner's vasectomy
				PLUS One method from this list:		PLUS One method from this list:
				Male condom		Male condom Diaphragm with spermicides Cervical cap with spermicides Estrogen and Progesterone oral contraceptives ("the pill") Estrogen and Progesterone transdermal patch Vaginal ring Progesterone injection

- Do not have unprotected sex. Talk to your doctor or pharmacist right away if you have unprotected sex or if you think your birth control has failed. Your doctor may tell you to use emergency birth control.
- Tell your doctor right away if you miss a menstrual period or think you may be pregnant for any reason.

If you are the parent or caregiver of a female child who started taking ambrisentan tablets before reaching puberty, you should check your child regularly to see if she is developing signs of puberty. Tell your doctor right away if you notice that she is developing breast buds or pubic hair. Your doctor should decide if your child has reached puberty. Your child may reach puberty before having her first menstrual period.

Females can only receive ambrisentan tablets through a restricted program called the Ambrisentan Risk Evaluation and Mitigation Strategy (REMS) program. If you are a female who can get pregnant, you must talk to your doctor, understand the benefits and risks of ambrisentan tablets, and agree to all of the instructions in the Ambrisentan REMS program.

Males can receive ambrisentan tablets without taking part in the Ambrisentan REMS program.

What are Ambrisentan Tablets?

- Ambrisentan tablets are a prescription medicine used to treat pulmonary arterial hypertension (PAH), which is high blood pressure in the arteries of your lungs.
- Ambrisentan tablets can improve your ability to exercise and it can help slow down the worsening of your physical condition and symptoms.
- It is not known if ambrisentan tablets are safe and effective in children.

Who should not take ambrisentan tablets?

Do not take ambrisentan tablets if:

- you are pregnant, plan to become pregnant, or become pregnant during treatment with ambrisentan tablets. Ambrisentan tablets can cause serious birth defects. (See "What is the most important information I should know about ambrisentan tablets?") Serious birth defects from ambrisentan tablets happen early in pregnancy.
- you have a condition called Idiopathic Pulmonary Fibrosis (IPF)
- What should I tell my doctor before taking ambrisentan tablets?
- Before you take ambrisentan tablets, tell your doctor if you:
 - have been told that you have a low red blood cell level (anemia)
 - have liver problems
 - have any other medical conditions

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Ambrisentan tablets and other medicines may affect each other, causing side effects. Do not start any new medicines until you check with your doctor.

Especially tell your doctor if you take the medicine cyclosporine (Gengraf, Neoral, Sandimmune). Your doctor may need to change your dose of ambrisentan tablets.

How should I take Ambrisentan Tablets?

- Ambrisentan tablets will be mailed to you by a certified pharmacy. Your doctor will give you complete details.
- Take ambrisentan tablets exactly as your doctor tells you to take it. Do not stop taking ambrisentan tablets unless your doctor tells you to stop.
- You can take ambrisentan tablets with or without food.
- Do not split, crush or chew ambrisentan tablets.
- It will be easier to remember to take ambrisentan tablets if you take it at the same time each day.
- If you take more than your regular dose of ambrisentan tablets, call your doctor right away.
- If you miss a dose, take it as soon as you remember that day. Take your next dose at the regular time. Do not take two doses at the same time to make up for a missed dose.

What should I avoid while taking Ambrisentan Tablets?

- Do not get pregnant while taking ambrisentan tablets. (See the serious birth defects section of the Medication Guide above called "What is the most important information I should know about ambrisentan tablets?") If you miss a menstrual period, or think you might be pregnant, call your doctor right away.
- It is not known if ambrisentan passes into your breast milk. You should not breastfeed if you are taking ambrisentan tablets. Talk to your doctor about the best way to feed your baby if you take ambrisentan tablets.

What are the possible side effects of ambrisentan tablets?

Ambrisentan tablets can cause serious side effects including:

- See "What is the most important information I should know about ambrisentan tablets?"
- Swelling all over the body (fluid retention) can happen within weeks after starting ambrisentan tablets. Tell your doctor right away if you have any unusual weight gain, tiredness, or trouble breathing while taking ambrisentan tablets. These may be symptoms of a serious health problem. You may need to be treated with medicine or need to go to the hospital.
- Decreased sperm count. Decreased sperm counts have happened in some men taking a medicine that is like ambrisentan tablets. A decreased sperm count may affect the ability to father a child. Tell your doctor if being able to have children is important to you.
- Low red blood cell levels (anemia) can happen during the first weeks after starting ambrisentan tablets. If this happens, you may need a blood transfusion. Your doctor will do blood tests to check your red blood cells before starting ambrisentan tablets. Your doctor may also do these tests during treatment with ambrisentan tablets.

The most common side effects of ambrisentan tablets include:

- swelling of hands, legs, ankles and feet (peripheral edema)
- stuffy nose (nasal congestion)
- inflamed nasal passages (sinusitis)
- hot flashes or getting red in the face (flushing)

Some medicines that are like ambrisentan tablets can cause liver problems. Tell your doctor if you get any of these symptoms of a liver problem while taking ambrisentan tablets:

- loss of appetite
- nausea or vomiting
- pain in the upper right stomach (abdominal) area
- yellowing of your skin or the whites of your eyes
- fever
- dark urine
- itching
- achiness
- generally do not feel well

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all of the possible side effects of ambrisentan tablets. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store ambrisentan tablets?

Store ambrisentan tablets at room temperature between 68°F to 77°F (20°C to 25°C), in the package it comes in.

Keep ambrisentan tablets and all medicines out of the reach of children.

General information about the safe and effective use of ambrisentan tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use ambrisentan tablets for a condition for which it was not prescribed. Do not give ambrisentan tablets to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about ambrisentan tablets. If you would like more information, ask your doctor. You can ask your doctor or pharmacist for information about ambrisentan tablets that is written for health professionals.

What are the ingredients in ambrisentan tablets?

Active ingredient: ambrisentan

Inactive Ingredients: croscarmellose sodium, lactose monohydrate, lecithin, magnesium stearate, microcrystalline cellulose, partially hydrolyzed polyvinyl alcohol, polyethylene glycol, povidone, talc and titanium dioxide. Additionally, 5 mg tablet contains: FD&C red#40 aluminum lake.